

Black-striped Wallaby (*Macropus dorsalis*)*

Milk	Age (days)	Foot (mm)	Head (mm)	Weight (g)	Feed (mL/day)	Notes
<0.4	60	24	33	35	11	
	70	28	37	45	13	Whiskers present
	80	33	41	60	16	
Transition	82	34	42	64	16	12mL <0.4 + 4ml 0.4
	85	36	43	71	16	8mL <0.4 + 8ml 0.4
	88	38	44	78	16	4mL <0.4 + 12ml 0.4
0.4	90	39	45	83	16	
	100	45	49	110	20	
	110	52	53	140	24	Eyes opening
	115	56	55	155	26	
Transition	117	58	55	162	26	20mL 0.4 + 6ml 0.6
	120	60	57	174	26	13mL 0.4 + 13ml 0.6
	123	63	58	186	26	6mL 0.4 + 20ml 0.6
0.6	125	64	58	195	26	
	130	69	60	220	29	
	140	79	64	280	34	Fine hair covering
Transition	142	81	65	296	34	26mL 0.6 + 8ml >0.7
	145	84	66	320	34	17mL 0.6 + 17ml >0.7
	148	87	67	344	34	8mL 0.6 + 26ml >0.7
>0.7	150	89	67	360	34	
	160	99	71	460	40	Emerging from pouch
	170	107	74	575	49	
	180	115	78	720	58	
	190	121	82	900	70	
	200	126	86	1100	80	
	210	131	90	1300	90	Fully out of pouch
Weaning	230	Growth rate now about 20g per day			80	Feed volume now depends on amount of solids consumed
	250	Not Valid			60	
	270				40	Gradually increase solid food and reduce milk intake
	290				20	
	310				0	

* Males may be significantly larger than females, especially once emerged from the pouch.