

## Southern Hairy-Nosed Wombat (*Lasiorhinus latifrons*)

Milk	Age (days)	Head (mm)	Weight (g)	Feed (mL/day)	Notes
<0.4	70	33	65	17	
	80	37	85	21	
	90	41	110	25	
	100	45	138	30	
	105	46	158	32	Incisors present, eyelashes growing
Transition	107	47	168	32	24mL <0.4 + 8ml 0.4
	110	48	185	32	16mL <0.4 + 16ml 0.4
	113	50	203	32	8mL <0.4 + 24ml 0.4
0.4	115	50	215	32	
	120	52	250	34	
	130	56	320	42	Eyes opening
	140	60	400	49	
	150	64	490	58	Fine fur visible
Transition	160	68	590	64	
	164	70	640	64	48mL 0.4 + 16ml 0.6
	168	71	690	64	32mL 0.4 + 32ml 0.6
>0.6	172	73	740	64	16mL 0.4 + 48ml 0.6
	175	74	780	64	
	180	76	850	70	
	190	80	1000	80	
	200	84	1160	90	Emerging from pouch
	210	88	1350	100	Thick Fur
	220	92	1550	110	
	230	96	1800	115	
	240	100	2100	130	
	250	103	2450	145	
Weaning	260	107	2800	155	
	270	111	3150	170	Fully emerged from pouch
	280	Growth rate now about 35g per day		160	Feed volume now depends on amount of solids consumed
	290			150	
	310			100	Gradually increase solid food and reduce milk intake
330	50				
350			0	Fully weaned (approx 6-7 kg)	